arXiv:hep-th/0409146v1 14 Sep 2004

Dynamical Generation of Yukawa Couplings in Intersecting D-brane Models

NORIAKI KITAZAWA
Department of Physics, Tokyo Metropolitan University,
1-1 Minami-Osawa, Hachioji, Tokyo 192-0397, Japan
E-mail: kitazawa@phys.metro-u.ac.jp

ABSTRACT

We propose a scenario to obtain non-trivial Yukawa coupling matrices for the quark-lepton mass generation in supersymmetric intersecting D-brane models in type IIA $\mathbf{T}^6/\mathbf{Z}_2 \times \mathbf{Z}_2$ orientifold. As an example, an explicit model is constructed in which all the four generations of quarks and leptons and two pairs of massless Higgs fields are composite. In this model non-trivial Yukawa interactions are obtained by the interplay between the string-level higher dimensional interactions among "preons" and the dynamics of the confinement of "preons".

1. Introduction

The understanding of masses and mixings of quarks and leptons is one of the most important problems in particle physics. It has been expected that the string theory can give a solution as well as the unified description of the fundamental interactions including gravity. Recent developments in the model building based on the intersecting D-branes (see Refs.[1,2,3] for essential idea) make the explicit and concrete discussions possible. Especially, the models with low-energy supersymmetry are interesting, because such models are constructed as stable solutions of the string theory (see Refs.[3,4,5,6,7,8,9,10,11,12,13] for explicit model buildings).

The structure of the Yukawa coupling matrices in intersecting D-brane models is discussed in Refs.[14,15,16,17,18]. The problem is that we typically have the factorized form of the Yukawa coupling matrices, $g_{ij} = a_i b_j$, or the diagonal Yukawa coupling matrices, if the origin of the generation is the multiple intersections of D-branes. Both structures can not give realistic quark or lepton masses and mixings. The efforts in the model building, including many Higgs doublets[17], for example, may solve the problem, but the origin of the generation may have to be reconsidered.

In this article, we propose another scenario to have generation structure. The generation may not be originated from the multiple intersections of D-branes, but the repetitive existence of many confining forces for "preons". For example, suppose that we have "preons" with some specific charge under the standard model gauge group which is appropriate to form one generation with the confining USp(2) gauge interaction. If such "preons" are realized in an intersecting D-brane model, and they belong to the fundamental representation of USp(6), then we have composite three generations by the decomposition of USp(6) → USp(2) × USp(2) × USp(2) due to the D-brane splitting. The difference of the positions of D-branes for each USp(2) gauge symmetry may affect the structure of Yukawa coupling matrices. In the following we sketch a model in which this idea is explicitly realized (see Ref.[11] for complete description). We also give an evidence that the resultant Yukawa coupling matrices in this scenario can be realistic.

2. The Model and Yukawa Coupling Matrices

The configuration of the intersecting D6-branes in type IIA $\mathbf{T}^2 \times \mathbf{T}^2 \times \mathbf{T}^2/\mathbf{Z}_2 \times \mathbf{Z}_2$ orientifold is given in Table 1.

Table 1: Configuration of intersecting D6-branes. All three tori are considered to be rectangular (untilted). Three D6-branes, D6_4, D6_5 and D6_6, are on top of some O6-planes. We also have orientifold image D-brane for each D-brane listed in this table.

D6-brane	winding number	multiplicity
D6_1	$[(1,-1),(1,1),(1,0)]$	4
D6_2	$[(1,1),(1,0),(1,-1)]$	$6+2$
D6_3	$[(1,0),(1,-1),(1,1)]$	$2+2$
D6_4	$[(1,0),(0,1),(0,-1)]$	12
D6_5	$[(0,1),(1,0),(0,-1)]$	8
D6_6	$[(0,1),(0,-1),(1,0)]$	12

Ramond-Ramond tadpoles are canceled out in this configuration, and four-dimensional $\mathcal{N}=1$ supersymmetry is realized under the condition of $\chi_1=\chi_2=\chi_3=\chi$, where $\chi_i = R_2^{(i)}/R_1^{(i)}$ and $R_{1,2}^{(i)}$ are radii for each three torus $i=1,2,3$. The D6_2-brane system consists of two parallel D6-branes with multiplicities six and two which are separated in the second torus in a consistent way with the orientifold projections. The D6_3-brane system consists of two parallel D6-branes with multiplicity two which are separated in the first torus in a consistent way with the orientifold projections. D6_1, D6_2 and D6_3 branes give gauge symmetries of U(2)$_L$ =SU(2)$_L$×U(1)$_L$, U(3)$_c$×U(1) =SU(3)$_c$×U(1)$_c$×U(1) and U(1)$_1$×U(1)$_2$, respectively. The hypercharge is defined as

$$\frac{Y}{2} = \frac{1}{2}\left(\frac{Q_c}{3} - Q\right) + \frac{1}{2}\left(Q_1 - Q_2\right), \tag{1}$$

where Q_c, Q, Q_1 and Q_2 are charges of U(1)$_c$, U(1), U(1)$_1$ and U(1)$_2$, respectively.

We break three USp(12)$_{\mathrm{D}6_4}$, USp(8)$_{\mathrm{D}6_5}$ and USp(12)$_{\mathrm{D}6_6}$ gauge symmetries to the factors of USp(2) gauge symmetries by appropriately configuring D6-branes of D6_4, D6_5 and D6_6 (see Ref.[11] for concrete configuration). The resultant gauge symmetries are respectively as follows.

$$\mathrm{USp}(12)_{\mathrm{D}6_4} \longrightarrow \bigotimes_{\alpha=1}^{6} \mathrm{USp}(2)_{\mathrm{D}6_4,\alpha} \tag{2}$$

$$\mathrm{USp}(8)_{\mathrm{D}6_5} \longrightarrow \bigotimes_{a=1}^{4} \mathrm{USp}(2)_{\mathrm{D}6_5,a} \tag{3}$$

$$\mathrm{USp}(12)_{\mathrm{D}6_6} \longrightarrow \bigotimes_{\alpha=1}^{6} \mathrm{USp}(2)_{\mathrm{D}6_6,\alpha} \tag{4}$$

All of these USp(2) gauge intersections can be naturally stronger than any other unitary gauge interactions. If we choose $\kappa_4 M_s \sim 1$ and $\chi \sim 0.1$, where $\kappa_4 = \sqrt{8\pi G_N}$ and $M_s = 1/\sqrt{\alpha'}$, the scales of dynamics of all USp(2) gauge interactions are of the order of

M_s, and the values of the standard model gauge coupling constants are reasonably of the order of 0.01 at the string scale.

A schematic picture of the configuration of intersecting D6-branes of this model is given in Figure 1.



Figure 1: Schematic picture of the configuration of intersecting D6-branes. This picture schematically shows the intersections of D6-branes in six-dimensional space, and the relative place of each D6-brane has no meaning. The number at the intersection point between $D6_a$ and $D6_b$ branes denotes intersection number I_{ab} with $a < b$.

"Preons" are localized on six apices of the hexagonal area. The sector of $D6_1$-$D6_2$-$D6_4$ intersections gives four generation of left-handed quarks and leptons as two-body bound states of "preons". There are six composite generations due to six USp(2) gauge interactions of $D6_4$-brane, and two anti-generations due to the twice intersection between $D6_1$-brane and $D6_2$-brane. Two of six generations become massive with two anti-generations through the Yukawa couplings among two "preons" and one anti-generation field associated with the triangular aria of this sector. The similar happens to the sector of $D6_2$-$D6_3$-$D6_6$ intersections which gives four generation of right-handed quarks and leptons. The sector of $D6_1$-$D6_3$-$D5_5$ intersections gives two pairs of massless Higgs doublets.

The hexagonal area in Figure 1 indicates the existence of six-point higher-dimensional interactions among "preons". Since all the quarks, leptons and Higgs fields are two-body bound state of "preons", the higher-dimensional interactions give Yukawa interactions after the confinement of USp(2) gauge interactions. The value and structure of Yukawa coupling matrices are determined by the positions of six intersections of D-branes in the compact space.

In Ref.[18] the possibility to obtain non-trivial Yukawa coupling matrices has been shown. The Yukawa coupling matrices of the heaviest two generations of up-type and down-type quarks in a certain condition of the sizes of three tori are given as follows.

$$g^u \sim \begin{pmatrix} 1 & 0 \\ \varepsilon_1^2\varepsilon_3 & \varepsilon_1\varepsilon_3 \end{pmatrix} + \mathcal{O}(\varepsilon_3^2), \qquad g^d \sim \begin{pmatrix} \varepsilon_1 & 0 \\ \varepsilon_1\varepsilon_3 & \varepsilon_3 \end{pmatrix} + \mathcal{O}(\varepsilon_3^2), \tag{5}$$

where $\varepsilon_i = \exp(-A_i/2\pi\alpha')$ and A_i is the 1/8 of the area of the i-th torus. Note that we can obtain Yukawa coupling of the order of unity, although it seems difficult that all six positions of intersections coincide. These Yukawa coupling matrices give mass ratio and Kobayashi-Maskawa mixing angles for heavy two generations in a certain assumption to the vacuum expectation values of Higgs doublet fields. The results are

$$\frac{m_{u,3}}{m_{u,4}} \sim \varepsilon_1\varepsilon_3, \qquad \frac{m_{d,3}}{m_{d,4}} \sim \frac{\varepsilon_3}{\varepsilon_1}, \qquad V_{34} \sim \varepsilon_3. \tag{6}$$

By taking $\varepsilon_1 \sim 0.5$ and $\varepsilon_3 \sim 0.01$, we obtain the values corresponding to $m_c/m_t \simeq 0.0038$, $m_s/m_b \simeq 0.025$, and $V_{cb} \simeq 0.04$.

It would be very interesting to explore more realistic models of the quark-lepton flavor in this framework.

3. Acknowledgements

The author thank T. Kobayashi, N. Maru and N. Okada, for useful discussions.

4. References

[1] R. Blumenhagen, L. Görlich, B. Körs and D. Lüst, JHEP **0010**, 006 (2000).
[2] G. Aldazabal, S. Franco, L.E. Ibáñez, R. Rabadán and A.M. Uranga, J. Math. Phys. 42, 3103 (2001); JHEP **0102**, 047 (2001).
[3] M. Cvetič, G. Shiu and A.M. Uranga, Phys. Rev. Lett. **87**, 201801 (2001); Nucl. Phys. **B615**, 3 (2001).
[4] R. Blumenhagen, L. Görlich and T. Ott, JHEP **0301**, 021 (2003).
[5] M. Cvetič, I. Papadimitriou and G. Shiu, Nucl. Phys. **B659**, 193 (2003).
[6] G. Honecker, Nucl. Phys. **B666**, 175 (2003).
[7] M. Larosa and G. Pradisi, Nucl. Phys. **B667**, 261 (2003).
[8] M. Cvetič and I. Papadimitriou, Phys. Rev. **D67**, 126006 (2003).
[9] N. Kitazawa, hep-th/0401096, to be published in Nucl. Phys. B.
[10] M. Cvetič, T. Li and T. Liu, hep-th/0403061.
[11] N. Kitazawa, hep-th/0403278.
[12] G. Honecker and T. Ott, hep-th/0404055.
[13] C. Kokorelis, hep-th/0406258.
[14] M. Cvetič, P. Langacker and G. Shiu, Nucl. Phys. **B642**, 139 (2002).
[15] D. Cremades L.E. Ibáñez and F. Marchesano, JHEP **0307**, 038 (2003).
[16] N. Chamoun, S. Khalil and E. Lashin, Phys. Rev. **D69**, 095011 (2004).
[17] S.A. Abel, O. Lebedev and J. Santiago, hep-ph/0312157.
[18] N. Kitazawa, T. Kobayashi, N. Maru and N. Okada, hep-th/0406115.